



Contribution ID: 104

Type: **Contributed talk**

Elliptic anisotropy measurement of the $f_0(980)$ hadron in proton-lead collisions and evidence for its quark-antiquark

Despite the $f_0(980)$ hadron having been discovered half a century ago, the question about its quark content has not been settled: it might be an ordinary quark-antiquark meson, a tetraquark exotic state, a kaon-antikaon molecule, or a quark-antiquark-gluon hybrid. We report strong evidence that the $f_0(980)$ state is an ordinary quark-antiquark meson, inferred from the scaling of elliptic anisotropies (v_2) with the number of constituent quarks (n_q), as empirically established using conventional hadrons in relativistic heavy ion collisions. The $f_0(980)$ state is reconstructed via its dominant $\pi\pi$ decay channel, in proton-lead collisions recorded by the CMS experiment at the LHC, and its v_2 is measured as a function of transverse momentum (p_T). It is found that the $n_q = 2$ (quark-antiquark state) hypothesis is favored over $n_q = 4$ (tetraquark or molecule states) by 7.7, 6.3, or 3.1 standard deviations in the $p_T < 10$, 8, or 6 GeV/c ranges, respectively, and over $n_q = 3$ (quark-antiquark-gluon hybrid state) by 3.5 standard deviations in the $p_T < 8$ GeV/c range. This result represents the first determination of the quark content of the $f_0(980)$ state, made possible by using a novel approach, and paves the way for similar studies of other exotic hadron candidates.

session

G. Heavy Ion Physics

Primary author: WANG, Fuqiang (Purdue University)

Presenter: WANG, Fuqiang (Purdue University)

Session Classification: G. Heavy Ion Physics