



Contribution ID: 114

Type: **Contributed talk**

## The impact of scale-invariant initial state for gaining insights into nucleus properties in ultra-central symmetric heavy-ion collisions.

*Thursday, 11 July 2024 15:20 (20 minutes)*

Heavy-ion collision experiments are a valuable tool for studying nuclear properties. Accurately modeling entropy production at the initial collision time and subsequent collective evolution is crucial to connect the nuclear structure to heavy-ion measurements. In this talk, we argue that, based on experimental data, it is reasonable to assume scale-invariance at the initial state, meaning the produced entropy scales similarly to the thickness function of the nuclei participants. This implies that the observables receive a small contribution from the entropy production process and more from nuclear properties in ultracentral symmetric heavy-ion collision. With this conclusion, we employ cluster expansion decomposition to study the ellipticity fluctuation and introduce a formula for it. The formula is common for all scale-invariant initial state models and depends on the one-body and two-body density of the colliding nuclei. We show that this result is compatible with initial state models that obey scale invariance, such as TrENTo, with various values for  $p$  and initial state models based on CGC. We also explore its implications for studying nuclear properties in the isobar ratio measurements.

### session

G. Heavy Ion Physics

**Primary author:** TAGHAVI, Seyed Farid (Technische Universitaet Muenchen (DE))

**Presenter:** TAGHAVI, Seyed Farid (Technische Universitaet Muenchen (DE))

**Session Classification:** G. Heavy Ion Physics