## QNP2024 - The 10th International Conference on Quarks and Nuclear Physics



Contribution ID: 105

Type: Leading contributed talk

## Beta-decay studies with the Total Absorption technique

Monday 8 July 2024 16:30 (25 minutes)

Conventional high-resolution techniques for  $\beta$ -decay spectroscopy utilize high-purity germanium detectors to measure individual  $\gamma$  rays emitted after  $\beta$  decay. However, these measurements are affected by the Pandemonium systematic error [1], resulting in many high-energy  $\gamma$  rays and a significant portion of the  $\beta$  strength being missed. The Total Absorption  $\gamma$ -ray Spectroscopy (TAGS) technique effectively addresses this issue [2, 3]. It relies on detecting the full energy of  $\gamma$  cascades following the decay, achieved through the use of large, high-efficiency scintillation crystals acting as calorimeters. This technique allows for a Pandemonium-free determination of the  $\beta$  strength, a fundamental quantity that depends on the underlying nuclear structure, thus the ideal tool for constraining theoretical models.

The TAGS technique has been successfully utilized in  $\beta$ -decay studies for many years, yielding important results relevant to nuclear structure, nuclear astrophysics, and applications in reactor and neutrino physics (see Ref. [3] for a recent review). In my talk, I will highlight some selected achievements. I will then introduce Experiment E891\_23 [4], which has been granted beam time at the GANIL laboratory in France. The experiment aims to measure the  $\beta$ -decay properties of several proton-rich nuclei in the Cr-Zn region, of significant interest for both nuclear structure [5, 6] and nuclear astrophysics. This experiment will be performed with a new-concept hybrid spectrometer, STARS, currently under development within the (NA)<sup>2</sup>STARS project [7]. STARS will be the world's first device to combine the large  $\gamma$  efficiency characteristic of TAGS calorimeters with the superior energy resolution and timing of LaBr<sub>3</sub>(Ce) crystals. This unique combination will enable unprecedented studies further away from nuclear stability.

[1] J.C. Hardy et al., Phys. Lett. B 71, 307 (1977).

[2] B. Rubio et al., J. Phys. G Nucl. Part. Phys. 31, S1477 (2005).

[3] A. Algora et al., Eur. Phys. J. A 57, 85 (2021).

[4] Experiment E891\_23: "Total Absorption Spectroscopy for Nuclear Structure and Nuclear Astrophysics" (spokespersons: M. Fallot, S.E.A. Orrigo, A.M. Sánchez Benítez), approved by the GANIL Program Advisory Committee, Nov. 2023.

[5] S.E.A. Orrigo et al., Phys. Rev. Lett. 112, 222501 (2014).

[6] S.E.A. Orrigo et al., Phys. Rev. C 93, 044336 (2016).

[7] Project (NA)<sup>2</sup>STARS: "Neutrinos, Applications and Nuclear Astrophysics with a Segmented Total Absorption with a higher Resolution Spectrometer" (spokesperson: M. Fallot), endorsed by the GANIL Scientific Council, Jan. 2023.

## session

I. Nuclear Structure and Reactions

Author: ORRIGO, Sonja (IFIC (CSIC-UV))

Presenter: ORRIGO, Sonja (IFIC (CSIC-UV))

Session Classification: I. Nuclear Structure and Reactions