QNP2024 - The 10th International Conference on Quarks and Nuclear Physics



Contribution ID: 174

Type: Contributed talk

Spectroscopic study of Kaonic nuclei using inclusive and exclusive ${}^{12}{\rm C}(K^-,p)$ reaction at J-PARC

Wednesday 10 July 2024 14:40 (20 minutes)

It has been argued that the formation of a molecular-like quasi-bound state, $\Lambda(1405)$, arises from the strong attraction between $\overline{K}N$ with isospin I = 0 channel. Furthermore, its strong attraction suggests the existence of kaonic nuclei in a three-body system, K^-pp , as well as in more complex many-body systems.

Some calculations suggest existence of various types of many-body systems involving Kaon and nucleons. However, apart from the observation of the K^-pp three-body system[1][2], there have been few clear confirmations for the existence of Kaonic nuclei. Therefore, it is crucial to systematically investigate Kaonic nuclei across a wide range of mass numbers not only for understanding $\overline{K}N$ interactions but also for obtaining information on the presence or absence of \overline{K} within the inner core region of neutron stars.

The J-PARC E05 experiment studied the interaction between \overline{K} and a residual nucleus by measuring an inclusive ${}^{12}C(K^-, p)$ spectrum. From the analysis, the depths of both the real and imaginary parts of the \overline{K} -nucleus optical potential were obtained to be (-80, 40) [MeV] by fitting the shallow bound region with a theoretically calculated spectrum[3]. In addition, a significant event excess was observed in the spectrum in the deeply bound region around 90 MeV in the \overline{K} binding energy. That excess fits well with a Breit-Wigner function whose binding energy is 90 MeV and width is 100 MeV, suggesting possible contribution from production of $\overline{K}NN$ or a bound state between an excited hyperon (Y) and a nucleus.

To investigate the reason behind the excess event, we conducted a new experiment called E42[4]. This experiment used the same reaction dibaryon search experiment. We used a GEM-based Time Projection Chamber, HypTPC, installed around the target to measure the decay-charged particles during this experiment. This measurement technique helped improve the signalto-noise ratio and allowed us to detect the Y nuclear state as a clear bump if it exists.

In this talk, we will summarize the E05 results and report preliminary results of the inclusive ${}^{12}C(K^-, p)$ spectrum and the exclusive analysis using the information on particle identification with HypTPC.

Reference

S. Ajimura *et al.* Phys. Let. B **789** (2019) 620-625
T. Yamaga *et al.* Phys. Rev. C **102**, 044002 (2020)
Y. Ichikawa *et al.*, PTEP **2020**, 123D01 (2020)
J.K. Ahn *et al.*, the proposal of J-PARC E42 experiment, Search for H-Dibaryon with a Large Acceptance Hyperon Spectrometer

session

J. Strange Nuclear Systems

Author: OURA, Fumiya (Tohoku University)

Co-authors: KANG, Byung-Min (Korea University); TAMURA, Hirokazu (Tohoku University, JAEA); SAKO, Hiroyuki (JAEA); AHN, Jung-Keun (Korea University); TANIDA, Kiyoshi (JAEA); KIM, Shin-Hyung (Kyungpook

National University); HAYAKAWA, Shuhei (Tohoku University); CHOI, Sung-Wook (Korea University); JUNG, Woo-Seung (Korea University); ICHIKAWA, Yudai (Tohoku University)

Presenter: OURA, Fumiya (Tohoku University)

Session Classification: J. Strange Nuclear Systems