

Search for Dark Matter annihilation with a combined analysis of dwarf spheroidal galaxies from

Fermi-LAT, HAWC, H.E.S.S., MAGIC, and VERITAS

7th Heidelberg International Symposium on High-Energy Gamma-Ray
Astronomy (γ -2022)
Barcelona, 7th July 2022


Daniel Kerszberg (IFAE-BIST), Céline Armand, Eric Charles, Mattia di Mauro, Chiara Giuri,
J. Patrick Harding, Tjark Miener, Emmanuel Moulin, Louise Oakes, Vincent Poireau, Elisa
Pueschel, Javier Rico, Lucia Rinchiuso, Daniel Salazar-Gallegos, Kirsan Tollefson, Benjamin Zitzer
for the Fermi-LAT, HAWC, H.E.S.S., MAGIC, and VERITAS collaborations

Dark Matter indirect searches

- **Looking for DM particles self-annihilating into SM particles**
- **Gamma-rays are not deflected by magnetic fields and trace back to original source**
 - Critical to identify the (physical) origin of the signal and study DM spatial distribution
- **Classical targets for gamma-ray experiments include among others:**
 - The Galactic Center (high DM content with high uncertainties)
 - Dwarf spheroidal galaxies (lower DM content with smaller uncertainties)
- **Here we will focus on dwarf spheroidal galaxies for which:**
 - The expected astrophysical gamma-ray emission is negligible
 - Large data sets have been already collected
 - Combining data from existing experiments allows to maximize the sensitivity to potential DM signals by increasing the statistics without requesting more observation time

Involved experiments

- Initiative by 5 gamma-ray experiments to combine their observations of dwarf galaxies:
 - Fermi-LAT
 - HAWC
 - H.E.S.S.
 - MAGIC
 - VERITAS



Fermi-LAT

- **Satellite in operation since 2008**
- **Energy range:
20 MeV - above 300 GeV**
- **Field of view ~20% of the sky**
- **Scan the whole sky every
~3 hours**



Fermi-LAT: in orbit at 550 km

HAWC

- **Array of water Cherenkov detectors in operation since 2013**
- **Energy range:
300 GeV - 100 TeV**
- **Field of view ~15% of the sky**



HAWC: Puebla, Mexico, 4100 m

H.E.S.S.


- **Array of five Cherenkov telescopes**
 - Phase I with 4 telescopes of 12 m diameter since 2003
 - Phase II with the addition of a telescope of 28 m diameter since 2012
- **Energy range:**
30 GeV - 100 TeV
- **Field of view of 5°**



H.E.S.S.: Khomas Highland, Namibia, 1800 m

MAGIC

- **MAGIC consists of two 17 m diameter Cherenkov telescopes**
 - First telescope since 2004
 - Second telescope since 2009
- **Energy range:
50 GeV - 50 TeV**
- **Field of view of $\sim 3.5^\circ$**



MAGIC: La Palma, Spain, 2200 m

VERITAS

- **Array of four 12 m diameter Cherenkov telescopes since 2007**
- **Energy range:
100 GeV - 30 TeV**
- **Field of view of 3.5°**



VERITAS: Arizona, USA, 1300 m

List of targets

- In this project we use a list of 20 dwarf galaxies for which individual collaborations already published results

- In total, 45 different data sets used

Source name	Fermi-LAT	HAWC	H.E.S.S., MAGIC, VERITAS		
	Exposure (10^{11} s m^2)	$ \Delta\theta $ ($^\circ$)	IACT	Zenith ($^\circ$)	Exposure (h)
Boötes I	2.6	4.5	VERITAS	15 – 30	14.0
Canes Venatici I	2.9	14.6	–	–	–
Canes Venatici II	2.9	15.3	–	–	–
Carina	3.1	–	H.E.S.S.	27 – 46	23.7
Coma Berenices	2.7	4.9	H.E.S.S.	47 – 49	11.4
			MAGIC	5 – 37	49.5
Draco	3.8	38.1	MAGIC	29 – 45	52.1
			VERITAS	25 – 40	49.8
Fornax	2.7	–	H.E.S.S.	11 – 25	6.8
Hercules	2.8	6.3	–	–	–
Leo I	2.4	6.7	–	–	–
Leo II	2.6	3.1	–	–	–
Leo IV	2.4	19.5	–	–	–
Leo V	2.4	–	–	–	–
Leo T	2.6	–	–	–	–
Sculptor	2.7	–	H.E.S.S.	10 – 46	11.8
Segue I	2.5	2.9	MAGIC	13 – 37	158.0
			VERITAS	15 – 35	92.0
Segue II	2.7	–	–	–	–
Sextans	2.4	20.6	–	–	–
Ursa Major I	3.4	32.9	–	–	–
Ursa Major II	4.0	44.1	MAGIC	35 – 45	94.8
Ursa Minor	4.1	–	VERITAS	35 – 45	60.4

Combined likelihood analysis

- **Expected gamma-ray flux from DM annihilation:**


$$\frac{d^2\Phi(\langle\sigma v\rangle, J)}{dEd\Omega} = \frac{1}{4\pi} \times \frac{\langle\sigma v\rangle}{2m_{\text{DM}}^2} \sum_f \text{BR}_f \frac{dN_f}{dE} \times \frac{dJ}{d\Omega}$$

- **Using as many common ingredients as possible:**
 - Common range of channels and DM masses:
 - From 5 GeV to 100 TeV using the DM spectra from Cirelli et al. [JCAP 1103:051, 2011]
 - Studied 7 annihilation channels in total
 - Same J-factor values and statistical uncertainties
- **Individual experiments shared likelihood profile for each dSph/channel/mass combination for a fixed value of the J-factor**
 - statistical uncertainties on the J-factor are taken into account (the J-factor being a **nuisance parameter** in the combined likelihood)

Combined likelihood analysis

- **Combined likelihood:**

$$\mathcal{L}(\langle\sigma v\rangle; \nu \mid \mathcal{D}_{\text{dSphs}}) = \prod_{l=1}^{N_{\text{dSphs}}} \mathcal{L}_{\text{dSph},l}(\langle\sigma v\rangle; J_l, \nu_l \mid \mathcal{D}_{l,\text{measured}}) \times \mathcal{J}_l(J_l \mid J_{l,\text{obs}}, \sigma_{\log J_l})$$




- **The combination was performed with two independent softwares:**

- glike: <https://doi.org/10.5281/zenodo.4028908>
- LklCombiner: <https://doi.org/10.5281/zenodo.4450884>

Combined likelihood analysis

- **Combined likelihood:**


$$\mathcal{L}(\langle\sigma v\rangle; \nu \mid \mathcal{D}_{\text{dSphs}}) = \prod_{l=1}^{N_{\text{dSphs}}} \mathcal{L}_{\text{dSph},l}(\langle\sigma v\rangle; J_l, \nu_l \mid \mathcal{D}_{l,\text{measured}}) \times \mathcal{J}_l(J_l \mid J_{l,\text{obs}}, \sigma_{\log J_l})$$



- **The combination was performed with two independent softwares:**

- glike: <https://doi.org/10.5281/zenodo.4028908>
- LklCombiner: <https://doi.org/10.5281/zenodo.4450884>

Combined likelihood analysis: an example for one dSph




- The total likelihood combines the likelihood of the 20 targets!

Uncertainty on the DM content

- **The J-factor estimation is the largest source of uncertainty in this analysis**
- **We used 2 sets of J-factors to compare the effect on the final results**
 - From A. Geringer-Sameth et al. [APJ 801:74, 2015]
 - From V. Bonnivard et al. [MNRAS 446:3002, 2015 and MNRAS 453:849, 2015]
- **Some dSphs are marginally affected but some are very affected**



Name	$\log_{10} J$ (\mathcal{GS} set) $\log_{10}(\text{GeV}^2\text{cm}^{-5}\text{sr})$	$\log_{10} J$ (\mathcal{B} set) $\log_{10}(\text{GeV}^2\text{cm}^{-5}\text{sr})$
Boötes I	$18.24^{+0.40}_{-0.37}$	$18.85^{+1.10}_{-0.61}$
Canes Venatici I	$17.44^{+0.37}_{-0.28}$	$17.63^{+0.50}_{-0.20}$
Canes Venatici II	$17.65^{+0.45}_{-0.43}$	$18.67^{+1.54}_{-0.97}$
Carina	$17.92^{+0.19}_{-0.11}$	$18.02^{+0.36}_{-0.15}$
Coma Berenices	$19.02^{+0.37}_{-0.41}$	$20.13^{+1.56}_{-1.08}$
Draco	$19.05^{+0.22}_{-0.21}$	$19.42^{+0.92}_{-0.47}$
Fornax	$17.84^{+0.11}_{-0.06}$	$17.85^{+0.11}_{-0.08}$
Hercules	$16.86^{+0.74}_{-0.68}$	$17.70^{+1.08}_{-0.73}$
Leo I	$17.84^{+0.20}_{-0.16}$	$17.93^{+0.65}_{-0.25}$
Leo II	$17.97^{+0.20}_{-0.18}$	$18.11^{+0.71}_{-0.25}$
Leo IV	$16.32^{+1.06}_{-1.70}$	$16.36^{+1.44}_{-1.65}$
Leo V	$16.37^{+0.94}_{-0.87}$	$16.30^{+1.33}_{-1.16}$
Leo T	$17.11^{+0.44}_{-0.39}$	$17.67^{+1.01}_{-0.56}$
Sculptor	$18.57^{+0.07}_{-0.05}$	$18.63^{+0.14}_{-0.08}$
Segue I	$19.36^{+0.32}_{-0.35}$	$17.52^{+2.54}_{-2.65}$
Segue II	$16.21^{+1.06}_{-0.98}$	$19.50^{+1.82}_{-1.48}$
Sextans	$17.92^{+0.35}_{-0.20}$	$18.04^{+0.50}_{-0.28}$
Ursa Major I	$17.87^{+0.56}_{-0.33}$	$18.84^{+0.97}_{-0.43}$
Ursa Major II	$19.42^{+0.44}_{-0.42}$	$20.60^{+1.46}_{-0.95}$
Ursa Minor	$18.95^{+0.26}_{-0.18}$	$19.08^{+0.21}_{-0.13}$

Combined limits



- **Combined limits are up to a factor 2-3 more constraining**

Comparison of the limits using two different sets of J-factors



- “Bonnivard” provides better limits than “Geringer-Sameth” by a factor 2-6

Conclusion

- **This analysis framework allows us to perform multi-instrument and multi-target analysis**
- **No significant DM signal was observed**
- **Combined limits range from 5 GeV to 100 TeV and improve individual limits up to a factor 2 to 3**
- **Using 2 different sets of J-factors we were able to study the systematic impact on the results:**
 - **limits can vary by a factor of 2 to 6**
 - **combining many targets allows to minimize the importance of single dSphs, particularly relevant when their J-factor is (very) uncertain**
- **Publication under preparation will include in total 7 annihilation channels**
- **Combination including other messengers such as neutrinos is possible!**