



Machine Learning approaches to Amplitude Analysis

Glòria Montaña

Universitat de Barcelona & ICCUB

 gmontana@fqa.ub.edu



Hadron and Nuclear Physics group workshop (HADNUCMAT)

University of Barcelona, January 26-28, 2026




UNIVERSITAT DE
BARCELONA



EXCELENCIA
MARÍA
DE MAEZTU
04/2025-03/2031


beatriu
de pinós bp





AI for Data Analysis and PreservaTion



Nuclear and Particle physics **Experiments:** our probe to Hadron Spectroscopy and QCD




GlueX experiment



Theorist's perspective: Scattering Amplitude

$$\langle f | \mathcal{S} | i \rangle = \delta_{fi} + i \langle f | \mathcal{T} | i \rangle$$


$$\langle p'_1 p'_2 | \mathcal{T} | p_1 p_2 \rangle \propto \mathcal{A}(s, t)$$



Theorist's perspective: Scattering Amplitude

$$\langle f | \mathcal{S} | i \rangle = \delta_{fi} + i \langle f | \mathcal{T} | i \rangle$$


$$\langle p'_1 p'_2 | \mathcal{T} | p_1 p_2 \rangle \propto \mathcal{A}(s, t)$$



Theorist's perspective: Scattering Amplitude

$$\langle f | \mathcal{S} | i \rangle = \delta_{fi} + i \langle f | \mathcal{T} | i \rangle$$

$$\langle p'_1 p'_2 | \mathcal{T} | p_1 p_2 \rangle \propto \mathcal{A}(s, t)$$



What does the blob really represent?


See talk by Vincent Mathieu

Properties of scattering amplitudes

Principles of scattering theory

- **Unitarity:** probability conservation (all possible outcomes must add up to 100%)
- **Analyticity:** causality (outcome cannot happen before interaction, smooth dependence on energy)
- **Crossing symmetry:** swapping particles and anti-particles (the mathematical description remains the same)

Discontinuity over the unitarity cut:




$$2 \text{Im } \mathcal{A}_{i \rightarrow f} \propto \sum_n \mathcal{A}_{i \rightarrow n} \mathcal{A}_{n \rightarrow f}^\dagger$$

Properties of scattering amplitudes

Principles of scattering theory

- **Unitarity:** probability conservation (all possible outcomes must add up to 100%)
- **Analyticity:** **causality** (outcome cannot happen before interaction, smooth dependence on energy)
- **Crossing symmetry:** swapping particles and anti-particles (the mathematical description remains the same)



Poles correspond to physical hadronic states


Branch points at the opening to N-particle scattering thresholds

Properties of scattering amplitudes

Principles of scattering theory

- **Unitarity:** probability conservation (all possible outcomes must add up to 100%)
- **Analyticity:** causality (outcome cannot happen before interaction, smooth dependence on energy)
- **Crossing symmetry:** swapping particles and **anti-particles** (the mathematical description remains the same)

$$A_{ab \rightarrow cd}(s, t, u) = A_{a\bar{c} \rightarrow \bar{b}d}(t, s, u)$$




Challenges in hadron spectroscopy

Experiments measure **physical observables** (e.g. cross section) in the **real axis** (physical region)


Most hadrons are **resonances** (poles of the scattering amplitude in the complex-energy plane)

Sometimes identified through “bumps” in the cross section




Pole parameters $\sqrt{s_R} = M_R - i\frac{\Gamma_R}{2}$

Challenges in hadron spectroscopy



- (Wide) **overlapping states**
- Partial wave analysis (PWA) techniques are complicated
- Pole positions from amplitude analyses rely on **theoretical assumptions**
- Assigning quantum numbers and internal structure (e.g. molecule vs tetraquark) is often **model dependent**


Challenges in hadron spectroscopy



- (Wide) **overlapping states**
- Partial wave analysis (PWA) techniques are complicated
- Pole positions from amplitude analyses rely on **theoretical assumptions**
- Assigning quantum numbers and internal structure (e.g. molecule vs tetraquark) is often **model dependent**

- Experimental data are inherently distorted by **detector effects** ▷ unfolded before extracting meaningful physics
- Traditional observables may not be adequate in **multi-dimensional space** (multi-particle final states)
- **Large datasets** ▷ difficult to manipulate/preserve


The A(i)DAPT collaborative effort



- Experimentalists
- Theorists
- ML experts


M. Battaglieri, Y. Li, A. Pilloni,
N. Sato, A. Szczepaniak,
G. Montana, G. Foti, L. Bibrzycki,
T. Alghamdi, T. Vittorini,
D. Glazier, D. Lersch, T. Reed,
M. Spreafico, and others

The A(i)DAPT collaborative effort



Develop **AI tools** to:

- Accurately fit data in multi-dimensional space
- Unfold detector effects
- Move from cross sections to scattering amplitudes




From cross section to scattering amplitude

Scattering amplitude \mathcal{A}

- underlying dynamics
- complex quantity: magnitude + phase

Cross section σ

- experimental observable
- related to $|\mathcal{A}|^2$, missing information about the phase




From cross section to scattering amplitude

Scattering amplitude \mathcal{A}

- underlying dynamics
- complex quantity: magnitude + phase


Cross section σ

- experimental observable
- related to $|\mathcal{A}|^2$, missing information about the phase




Use physics-informed **generative AI**

From cross section to scattering amplitude with physics-informed generative AI



From cross section to scattering amplitude with physics-informed generative AI



Principles of scattering theory


From cross section to scattering amplitude with physics-informed generative AI

Goal: physically consistent amplitude reconstruction

- Single underlying physics (no model uncertainty)
- Noisy replicas \approx pseudo-experiments
- Strong physical constraints (unitarity)

Use ML as a regulator of the inverse problem

Generative Adversarial Network (GAN)



$\pi^+\pi^-$ scattering

Simplest case: elastic $2 \rightarrow 2$, spinless particles \triangleright 2 variables (s, t)

- Dominated by $f_0(500)$ and $\rho(770)$ resonances



Model of the amplitude with 2 partial waves:

$$\mathcal{A}(s, \cos \theta) = \sum_{\ell=0}^n (2\ell + 1) f_{\ell}(s) P_{\ell}(\cos \theta)$$

$$\mathcal{A}(s, \cos \theta) = f_0(s) + 3f_1(s) \cos \theta$$

Breit-Wigner type


$$f_{\ell} = \frac{m_{\ell} \Gamma_{\ell}}{m_{\ell}^2 - s - im_{\ell} \Gamma_{\ell}}$$



$\pi^+\pi^-$ scattering

Differential cross section


$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \frac{1}{s} |\mathcal{A}(s, \cos\theta)|^2$$



Unitarity relation


- For the partial waves $f_\ell(s) = \frac{1}{2} \int_{-1}^{+1} dz P_\ell(z) \mathcal{A}(s, z)$

$$\text{Im } f_\ell(s) = |f_\ell(s)|^2$$



Amplitudes and cross sections as images (discretized functions)

GAN with physics constraints




$$\mathcal{L} = \mathcal{L}_{\text{GAN}} + \lambda_u \mathcal{L}_u + \lambda_\delta \mathcal{L}_\delta$$

GAN with physics constraints



Training data: differential cross section from the model

- discretized 64×64 , $s \in [(2m_\pi)^2, 1 \text{ GeV}^2]$, $\cos \theta \in [-1, 1]$
- normalized
- noisy replicas \triangleright pseudo-data



GAN with physics constraints

With $\text{std}=0.01$, convergence to the ground truth for $\sim 30\%$ of trainings



GAN generated ("fake")

With $\text{std}=0.05$, hard to converge (unitarity is not satisfied exactly)

Next steps

- Generate $\text{Im } \mathcal{A}$ and use unitarity to reconstruct $\text{Re } \mathcal{A}$ instead of a constraint in the loss function
- Uncertainty quantification
- Explore other generative models

Next steps

- Generate $\text{Im } \mathcal{A}$ and use unitarity to reconstruct $\text{Re } \mathcal{A}$ instead of a constraint in the loss function
- Uncertainty quantification
- Explore other generative models



After the proof of concept with $\pi^+\pi^-$ scattering...

- Coupled channels $\pi^+\pi^-$, K^+K^-
- Spin
- $2 \rightarrow 3$ processes
- Meson photoproduction

Back-up slides

Conditional diffusion model

Training data:


- ▷ sample amplitudes and cross sections from 2BW model with parameters

$$m_0, m_1 \in [0.1, 1.0] \text{ GeV}$$

$$\Gamma_0, \Gamma_1 \in [0.01, 0.7] \text{ GeV}$$

- ▷ normalized
- ▷ with additional Gaussian noise to cross section (mimic experimental data)


Satisfy unitarity by construction




...

Conditional diffusion model

Forward diffusion process




Training



Conditional diffusion model

▷ With $\text{std}=0.01$

model (ground truth) Mean over 500 generated samples



Conditional diffusion model

▷ With std=0.05

model (ground truth) Mean over 500 generated samples

